

## Product Information

# HumanKine™ Macrophage-Colony Stimulating Factor, Human Recombinant, expressed in HEK 293 cells

**H6916**

Storage Temperature –20 °C

Synonyms: M-CSF, CSF-1

## Product Description

HumanKine™ M-CSF is expressed as a glycosylated 35-40 kDa disulfide linked homodimer in human 293 cells. Production in human 293 cells offers authentic glycosylation. Glycosylation contributes to stability in cell growth media and other applications.

Four distinct colony-stimulating factors (CSFs) that promote survival, proliferation, and differentiation of bone marrow precursor cells have been well characterized: granulocyte macrophage-CSF (GM-CSF), granulocyte-CSF (G-CSF), macrophage-CSF (M-CSF), and Interleukin-3 (IL-3, Multi-CSF).<sup>1,2</sup> Both GM-CSF and IL-3 are multipotential growth factors, stimulating proliferation of progenitor cells from more than one hematopoietic lineage. In contrast, G-CSF and M-CSF are lineage restricted hematopoietic growth factors, stimulating final mitotic divisions and the terminal cellular maturation of the partially differentiated hematopoietic progenitors.

Macrophage CSF (M-CSF) is produced by monocytes, fibroblasts, and endothelial cells. M-CSF stimulates the formation of macrophage colonies,<sup>3</sup> enhances antibody-dependent, cell-mediated cytotoxicity by monocytes and macrophages,<sup>4</sup> and inhibits bone resorption by osteoclasts.<sup>5</sup>

This product is lyophilized from a PBS solution.

EC<sub>50</sub>: ≤ 8.0 ng/mL

The specific activity was determined by the dose dependent stimulation of the proliferation of murine M-NFS-60 cells (mouse myeloid leukemia indicator cell line).

Purity: ≥ 95% (SDS-PAGE)

Endotoxin level: ≤ 1 EU/mg

## Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Preparation Instructions

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile PBS containing 0.1% endotoxin-free recombinant human serum albumin.

## Storage/Stability

Store the product at –20 °C.

Upon reconstitution, the cytokine can be stored at 2-8 °C for short term only, or at –20 °C to –80 °C in aliquots for long term. Avoid repeated freeze-thaw cycles.

---

## References

1. Mazur, E.M., and Cohen, J.L., Clin. Pharmacol. Ther., 46, 250 (1989).
2. Morstyn, G., and Burgess, A.W., Cancer Res., 48, 5624 (1988).
3. Metcalf, D., Blood, 67, 257 (1986).
4. Mufson, R.A., et al., Cell. Immunol., 119, 182 (1989).
5. Hattersley, G., et al., J. Cell Physiol., 137, 199 (1988).

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

## Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

## Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

## Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck operates  
as MilliporeSigma in the U.S. and Canada.

Merck, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2010-2025 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.  
H6916pis REV 01/252

